NH Lab Services

Blood Collection Order of Draw for Common Tests

Order of Draw	TEST		Tube Type	Tube Top Colour	Comment	# Inversions			
Tubes must be collected in the order indicated to prevent contamination or interference from a preceding tube to a following tube.									
1.	Non-Additive Used as discard tube prior to coagulation tube or prior to all tubes when using a butterfly needle		Clear Brown	Clear Top		DISCARD			
2.	Hematology Coagulation INR PTT		Coag/ Sodium Citrate	Light Blue	Volume is critical Draw to fill line	5			
3.	CRP Ferritin IgG, IgA, IgM Protein Serum free light chains Thyroglobulin (fasting) Thyroglobulin antibody TSH Vit B12 EP HIV Hepatitis	Tumour Markers AFP B2M CA 15-3 CA 125 CA 19-9 CEA Total hCG Total PSA Free PSA SCC Testosterone B2M	Serum Separator Tube	Gold	Gel separator Allow to clot 30 min	5			



Order of Draw	TEST	Tube Type	Tube Top Colour	Comment	# Inversions
4.	Albumin Glucose ALK (ALP) (fasting & ALT random) Amylase Iron AST LD BUN Magnesium *Bilirubin* Phosphate (Total & Direct) Potassium	Separator Tubes	PST Light Green Li Heparin (certain labs only)	PST tube - no clot time RST tube - clot time 5 mins	8
	Calcium Sodium Chloride TIBC Cholesterol (fasting) Triglycerides Creatinine (fasting) CK Uric Acid CO2 Urea GFR GGT (y-GT) *Protect from Light*		Gold (SST) or Orange (RST) (certain labs only)	SST tube - clot time 30 mins	5
5.	Hematology CBC/Differential/ Platelets Reticulocyte Count (order CBC with Retic count)	CBC	Lavender EDTA	Small purple top	8
6.	Crossmatch X 2 tubes min, x 3 is optimal	Crossmatch	Lavender EDTA	Large purple top	8
7.	Hematology ESR Note: written indication for ESR must be provided on requisition	ESR	Black Buffered	Volume is critical. Draw to fill line	8

